

INVARIANT BILINEAR FORMS ON A VERTEX ALGEBRA AND THE RADICAL

MICHAEL ROITMAN

INTRODUCTION

Invariant bilinear forms on vertex algebras have been around for quite some time now. They were mentioned by Borchers in [1] and were used in many early works on vertex algebras, especially in relation with the vertex algebras associated with lattices [2, 7, 11]. The first systematic study of invariant forms on vertex algebras is due to Frenkel, Huang and Lepowsky [6]. This theory was developed further by Li [12].

However, these authors imposed certain assumptions on their vertex algebras which are too restrictive for the applications we have in mind. Specifically, this paper is motivated by the study of vertex algebras of OZ type generated by their Griess subalgebras [9, 17]. We show that all the results of Li [12] hold in the greatest possible generality, basically, as long as the definitions make sense. We construct a linear space that parameterizes all bilinear forms on a given vertex algebra with values in an arbitrary linear space, and also prove that every invariant bilinear form on a vertex algebra is symmetric. Our methods, however, are very different from the methods used in [6] and [12].

As an application, we introduce a very useful notion of radical of a vertex algebra. It is equal to the radical of a certain canonical invariant bilinear form. We prove that a radical-free vertex algebra A is non-negatively graded, and its component A_0 of degree 0 is a commutative associative algebra, so that all structural maps and operations on A are A_0 -linear. We also show that in this case A is simple if and only if A_0 is a field.

1. DEFINITIONS AND NOTATIONS

Here we fix the notations and give some minimal definitions. For more details on vertex algebras the reader can refer to the books [5, 7, 11]. All spaces and algebras are considered over a field $\mathbb{k}$ of characteristic 0. We use the following notation for divided powers:

$$x^{(n)} = \begin{cases} (x^n)/n! & \text{if } n \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

1.1. Definition of vertex algebras.

Definition 1. A vertex algebra is a linear space A equipped with a family of bilinear products $a \otimes b \mapsto a(n)b$, indexed by integer parameter n , and with an element $\mathbb{1} \in A$, called the unit, satisfying the identities (i)–(iv) below. Let $D : A \rightarrow A$ be the map defined by $Da = a(-2)\mathbb{1}$. Then

- (i) $a(n)b = 0$ for $n \gg 0$,
- (ii) $\mathbb{1}(n)a = \delta_{n,-1}a$ and $a(n)\mathbb{1} = D^{(-n-1)}a$,
- (iii) $D(a(n)b) = (Da)(n)b + a(n)(Db)$ and $(Da)(n)b = -na(n-1)b$,
- (iv) $a(m)(b(n)c) - b(n)(a(m)c) = \sum_{s \geq 0} \binom{m}{s} (a(s)b)(m+n-s)c$

for all $a, b, c \in A$ and $m, n \in \mathbb{Z}$.

Date: February 8, 2008.

1991 Mathematics Subject Classification. 17B69.

This is not the only known definition of vertex algebras. Often the axioms are formulated in terms of the *left adjoint action map* (a.k.a. *state-field correspondence*) $Y : A \rightarrow A[[z, z^{-1}]]$ defined by $Y(a)(z) = \sum_{n \in \mathbb{Z}} a(n) z^{-n-1}$, where $a(n) : A \rightarrow A$ is the operator given by $b \mapsto a(n)b$. The most important property of these maps is that they are *local*: for any $a, b \in A$ there is N such that

$$[Y(a)(w), Y(b)(z)] (w - z)^N = 0.$$

The minimal $N = N(a, b)$ for which this identity holds is called *the locality index* of a and b . In fact, $N(a, b) = \min \{n \in \mathbb{Z} \mid a(m)b = 0 \ \forall m \geq n\}$. From (iii) we get $Y(Da)(z) = \partial_z Y(a)(z)$, which in terms of coefficients means

$$(1) \quad [D, a(m)] = -m a(m-1).$$

Among other identities that hold in vertex algebras are the associativity

$$(2) \quad (a(n)b)(m)c = \sum_{s \geq 0} (-1)^s \binom{n}{s} a(n-s)(b(m+s)c) - \sum_{s \leq n} (-1)^s \binom{n}{n-s} b(m+s)(a(n-s)c),$$

and the quasi-symmetry

$$(3) \quad a(n)b = - \sum_{i \geq 0} (-1)^{n+i} D^{(i)}(b(n+i)a).$$

For $n \geq 0$ the associativity identity (2) simplifies to

$$(a(n)b)(m)c = \sum_{s=0}^n (-1)^s \binom{n}{s} [a(n-s), b(m+s)] c,$$

which is just another form of the identity (iv) of Definition 1.

A vertex algebra A is called *graded* (by the integers) if $A = \bigoplus_{d \in \mathbb{Z}} A_d$ is a graded space, $A_i(n)A_j \subseteq A_{i+j-n-1}$ and $\mathbb{1} \in A_0$.

It is often required that a vertex algebra A is graded and A_2 contains a special element ω such that $\omega(0) = D$, $\omega(1)|_{A_d} = d$ and the coefficients $\omega(n)$ generate a representation of the Virasoro Lie algebra:

$$[\omega(m), \omega(n)] = (m-n)\omega(m+n-1) + \delta_{m+n,2} \frac{1}{2} \binom{m-1}{3} c$$

for some constant $c \in \mathbb{k}$ called the *central charge* of A . In this case A is called *conformal vertex algebra* or *vertex operator algebra*, especially when $\dim A_d < \infty$.

1.2. The action of sl_2 . In order to work with bilinear forms, we need to deal with vertex algebras equipped with certain additional structure. First of all we will assume that our vertex algebra $A = \bigoplus_{d \in \mathbb{Z}} A_d$ is graded. We will also need a locally nilpotent operator $D^* : A \rightarrow A$ of degree -1 , such that $D^*\mathbb{1} = 0$ and

$$(4) \quad [D^*, a(m)] = (2d - m - 2)a(m+1) + (D^*a)(m)$$

for every $a \in A_d$. Let $\delta : A \rightarrow A$ be the grading derivation, defined by $\delta|_{A_d} = d$. It is easy to compute that

$$(5) \quad [D^*, D] = 2\delta, \quad [\delta, D] = D, \quad [\delta, D^*] = -D^*,$$

so that D^*, D and δ span a copy of sl_2 .

An element $a \in A$ such that $D^*a = 0$ is called *minimal*. It is easy to see that if A is generated by minimal elements, then any operator $D^* : A \rightarrow A$ satisfying (4) must be locally nilpotent. For any $a \in A$ define $\text{ord } a = \min \{k \in \mathbb{Z}_+ \mid (D^*)^{k+1}a = 0\}$, so that $\text{ord } a = 0$ for a minimal a .

Vertex algebras with an action of sl_2 as above were called *quasi-vertex operator algebras* in [6] and minimal elements are sometimes called *quasi-primary*.

If A has a Virasoro element ω , then we always have $D = \omega(0)$ and $\delta = \omega(1)$, and we can take $D^* = \omega(2)$.

Throughout this paper we will always assume that all vertex algebras have an sl_2 -structure, and all ideals, homomorphisms, etc must agree with this structure. For example, we call a vertex algebra *simple* if it does not have D^* -invariant ideals. Note that any D^* -invariant ideal is homogeneous, because it must be stable under the grading derivation δ .

In the sequel we will need the following easy property of sl_2 -modules of the above type:

Proposition 1. *Let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a graded module over the Lie algebra $sl_2 = \mathbb{k}D + \mathbb{k}\delta + \mathbb{k}D^*$, where $\deg D = 1$, $\deg D^* = -1$, D^* is locally nilpotent, $\delta|_{M_d} = d$ and the commutation relations (5) hold. Then $M_d = (D^*)^{d+1}M_1$ for any $d < 0$.*

Proof. Let us first prove that $M_d = D^*M_{d+1}$ for $d < 0$. Take some element $a \in M_d$ of $\text{ord } a = k$. Set

$$b = D^*Da - (k+1)(2d-k)a = DD^*a - k(2d-k-1)a \in M_d.$$

If a is minimal, then $b = 0$. Otherwise, using that $(D^*)^{k+1}Da = (k+1)(2d-k)(D^*)^ka$, we get $(D^*)^kb = 0$, therefore $\text{ord } b < k$. By induction we can assume that $b \in D^*M_{d+1}$, and hence also $a \in D^*M_{d+1}$, because $(k+1)(2d-k) \neq 0$ for $d < 0$ and $k \geq 0$.

Now the proposition follows from the following claim:

$$M_0 = D^*M_1 + \text{Ker } D^*.$$

Take an element $a \in M_0$ of $\text{ord } a = k$. We will prove by induction on k that $a \in D^*M_1 + \text{Ker } D^*$. If $k = 0$ then $a \in \text{Ker } D^*$. Otherwise, take as before $b = D^*Da + k(k+1)a$ so that $\text{ord } b < k$. By induction, $b \in D^*M_1 + \text{Ker } D^*$, and therefore also $a \in D^*M_1 + \text{Ker } D^*$, since $k(k+1) \neq 0$ for $k > 0$. $\square$

It follows from Proposition 1 that $DM_{-1} \subset D^*M_1$. Indeed, if $a \in M_{-1}$, then $a = D^*b$ for some $b \in M_0$, and we have $Da = DD^*b = D^*Db$.

For the case when $M = A$ is a vertex algebra we will obtain that $A_d = D^*A_{d+1}$ as a corollary of Lemma 1 in §3.

Remark. It is possible (but not easy) to show that if a vertex algebra A is generated by minimal elements, then $A_d = D^*A_{d+1}$ for all $d \neq 0$. In contrast to that, the space A_0/D^*A_1 is non-zero in most interesting cases. We shall see in §3 that its dual space $\text{Hom}(A_0/D^*A_1, \mathbb{k})$ parameterizes the invariant bilinear forms on A .

1.3. The universal enveloping algebra. For any vertex algebra A we can construct a Lie algebra $L = \text{Coeff } A$ in the following way [1, 11, 13, 15]. Consider the linear space $\mathbb{k}[t, t^{-1}] \otimes A$, where t is a formal variable. Denote $a(n) = a \otimes t^n$ for $n \in \mathbb{Z}$. As a linear space, L the quotient of $\mathbb{k}[t, t^{-1}] \otimes A$ by the subspace spanned by the relations $(Da)(n) = -na(n-1)$. The brackets are given by

$$(6) \quad [a(m), b(n)] = \sum_{i \geq 0} \binom{m}{i} (a(i)b)(m+n-i),$$

which is precisely the identity (iv) of Definition 1. The spaces $L_{\pm} = \text{Span}\{a(n) \mid n \gtrless 0\} \subset L$ are Lie subalgebras of L and we have $L = L_- \oplus L_+$.

Remark. The construction of L makes use of only the products (n) for $n \geq 0$ and the map D . This means that it works for a more generic algebraic structure, known as *conformal algebra* [11].

The formulas (1) and (4) define derivations $D : L \rightarrow L$ and $D^* : L \rightarrow L$ so we get an action of sl_2 on L by derivations. Denote by $\widehat{L} = L \ltimes sl_2$ the corresponding semi-direct product.

Now assume that the vertex algebra A is $\mathbb{Z}$ -graded. Then so is the Lie algebra $\widehat{L} = \widehat{\text{Coeff } A}$ and its universal enveloping algebra $U = U(\widehat{L})$. The *Frenkel-Zhu topology* [8] on a homogeneous component U_d is defined by setting the neighborhoods of 0 to be the spaces $U_d^k = \sum_{i \leq k} U_{d-i}U_i$, so that

$$\dots \subset U_d^{k-1} \subset U_d^k \subset U_d^{k+1} \subset \dots \subset U_d, \quad \bigcap_{k \in \mathbb{Z}} U_d^k = 0, \quad \bigcup_{k \in \mathbb{Z}} U_d^k = U_d.$$

Let $\overline{U}(\widehat{L}) = \bigoplus_{d \in \mathbb{Z}} \overline{U}_d$ be the completion of U in this topology.

Definition 2. [8] The universal enveloping algebra $U(A)$ of a graded vertex algebra A is the quotient of $\overline{U}(\widehat{\text{Coeff } A})$ modulo the ideal generated by the relations

$$(a(n)b)(m) = \sum_{s \geq 0} (-1)^s \binom{n}{s} a(n-s)b(m+s) - \sum_{s \leq n} (-1)^s \binom{n}{n-s} b(m+s)a(n-s)$$

for $a, b \in A$ and $m, n \in \mathbb{Z}$.

Note that the relations above are simply the associativity identity (2). It is enough to impose these relations only for $n < 0$, because for $n \geq 0$ they are equivalent to (6).

A graded space M is a module over a vertex algebra A if it is a $U(A)$ -module that agrees with the Frenkel-Zhu topology on $U(A)$, i.e. a convergent sequence of elements from $U(A)$ makes sense as an operator on M .

2. THE INVOLUTION ON THE UNIVERSAL ENVELOPING ALGEBRA

Let A be a graded vertex algebra with sl_2 -structure, as in §1.2.

Proposition 2. *Set*

$$(7) \quad a(m)^* = (-1)^{\deg a} \sum_{i \geq 0} (D^{*(i)} a) (2 \deg a - m - 2 - i)$$

for a homogeneous $a \in A$ and $m \in \mathbb{Z}$, and also $D^{**} = D$ and $\delta^* = \delta$. Then there is a continuous anti-involution on the enveloping algebra $U(A)$ defined by $x \mapsto x^*$.

Note that $U(A)_d^* \subseteq U(A)_{-d}$.

Remark. Let M be a module over A . Assume that Proposition 2 is true. Then the graded dual space $M' = \bigoplus_{d \in \mathbb{Z}} \text{Hom}(M_d, \mathbb{k})$ also becomes an A -module, called the *contragredient module* of M , where the action of A is given by $(a(n)f)(v) = f(a(n)^*v)$, $(Df)(v) = f(D^*v)$ and $(D^*f)(v) = f(Dv)$ for $a \in A$, $f \in M'$, $v \in M$ and $n \in \mathbb{Z}$. Contragredient modules were introduced in [6] without the use of Proposition 2. In fact, it is easy to show that Proposition 2 is equivalent to the existence of the contragredient module. For the sake of completeness though, we present an independent proof of Proposition 2, that avoids the use of formal calculus.

Proof of Proposition 2. It is straightforward to check that $a(m)^{**} = a(m)$. This is particularly easy to see when $D^*a = 0$, and that already settles the case when A is generated by minimal elements. The general case is not much more difficult.

Let us show that (7) defines an anti-endomorphism of $U(A)$. We do this in three steps.

Step 1. We show first that (7) defines an anti-endomorphism of the coefficient Lie algebra $\widehat{L} = \text{Coeff}(A) \ltimes sl_2$. We have to check that

$$(8) \quad \begin{aligned} [b(n)^*, a(m)^*] &= \sum_{s \geq 0} \binom{m}{s} (a(s)b)(m+n-s)^*, \\ [a(m)^*, D^*] &= -m a(m-1)^* \end{aligned}$$

for a, b running over a set of generators of A and $m, n \in \mathbb{Z}$. Let us check this when a and b are minimal, thus proving (8) for the case when A is generated by the minimal elements. The general case is not much different.

Let $\deg a = d$, $\deg b = e$ and $D^*a = D^*b = 0$. Then we get

$$\begin{aligned} [b(n)^*, a(m)^*] &= -[(-1)^d a(2d-m-2), (-1)^e b(2e-n-2)] \\ &= (-1)^{d+e+1} \sum_{s \geq 0} \binom{2d-m-2}{s} (a(s)b)(2d+2e-m-n-s-4) \end{aligned}$$

On the other hand, using that

$$D^{*(i)}(a(s)b) = \binom{2d-s-2}{i} a(s+i)b,$$

we have

$$\begin{aligned} &\sum_{s \geq 0} \binom{m}{s} (a(s)b)(m+n-s)^* \\ &= \sum_{i, s \geq 0} (-1)^{d+e+s+1} \binom{m}{s} \binom{2d-s-2}{i} (a(s+i)b)(2d+2e-m-n-i-s-4) \end{aligned}$$

$$= (-1)^{d+e+1} \sum_{j \geq 0} \sum_{s=0}^j (-1)^s \binom{m}{s} \binom{2d-2-s}{j-s} (a(j)b)(2d+2e-m-n-j-4),$$

and the first equality of (8) follows from a binomial identity. Similarly, by (4) and (7),

$$[D^*, a(m)^*] = (-1)^d [D^*, a(2d-m-2)] = (-1)^d m a(2d-m-1) = m a(m-1)^*,$$

and this proves the second equality of (8).

Step 2. Assume that we have two vertex algebras A and B , both with sl_2 -structures as above. Assume also that for B the proposition is known, that is, formula (7) defines an anti-involution of $U(B)$, and suppose we have a surjective homomorphism $\varphi : B \rightarrow A$ of vertex algebras, which preserves the sl_2 -structure. Then the proposition also holds for A .

Indeed, we can extend φ to a homomorphism $U(\varphi) : U(B) \rightarrow U(A)$ of enveloping algebras. Since φ is surjective, so is $U(\varphi)$, and therefore $U(A) = U(B)/\text{Ker } U(\varphi)$. It is easy to see that $\text{Ker } U(\varphi)$ is the ideal of $U(B)$ generated by the coefficients $a(m)$ for all $a \in \text{Ker } \varphi \subset B$ and $m \in \mathbb{Z}$. But if $a \in \text{Ker } \varphi$, then also $D^*a \in \text{Ker } \varphi$, therefore $a(n)^* \in \text{Ker } U(\varphi)$, and it follows that $(\text{Ker } U(\varphi))^* \subseteq \text{Ker } U(\varphi)$.

Step 3. Consider the Verma $\widehat{L}$ -module $B = U(\widehat{L}) \otimes_{U(L_+) \ltimes sl_2} \mathbb{k}\mathbb{1}$ generated by a vector $\mathbb{1}$ such that $L_+\mathbb{1} = 0$ and also $D\mathbb{1} = D^*\mathbb{1} = 0$. Then B has a structure of vertex algebra such that $U(B)$ is the completion in the Frenkel-Zhu topology of the universal enveloping algebra $U = U(\widehat{L})$ of $\widehat{L}$. The action of D^* on B is given by (4), therefore Step 1 implies that (7) defines an anti-involution of U .

Now we show that this anti-involution on U is continuous in the Frenkel-Zhu topology, and therefore it can be extended to $U(B) = \overline{U}$. Indeed, let $u_1, u_2, \dots \in U_d$ be a convergent sequence. This means that for any $k \in \mathbb{Z}$ all elements u_n with sufficiently large n belong to $U_{d-i}U_i$ for some $i \leq k+d$. But then $u_n^* \in (U_{d-i}U_i)^* \subseteq U_{-i}U_{i-d} \subseteq U_{-d}^k$, therefore the sequence $\{u_n^*\}$ also converges in the Frenkel-Zhu topology.

Since A is also an $\widehat{L}$ -module generated by an element $\mathbb{1}$ such that $L_+\mathbb{1} = D\mathbb{1} = D^*\mathbb{1} = 0$, there is a unique $\widehat{L}$ -module homomorphism $\varphi : B \rightarrow A$ such that $\varphi(\mathbb{1}) = \mathbb{1}$. It is well known [3, 14, 15] that φ must be a vertex algebra homomorphism preserving the sl_2 -module structures. Therefore, Step 2 implies that (7) gives an involution of $U(A)$. $\square$

3. INVARIANT BILINEAR FORMS

Let as before A be a vertex algebra with sl_2 -structure. Let V be a vector space over $\mathbb{k}$.

Definition 3. A V -valued bilinear form $\langle \cdot | \cdot \rangle$ on A is called *invariant* if

$$\langle a(m)b | c \rangle = \langle b | a(m)^*c \rangle \quad \text{and} \quad \langle Da | b \rangle = \langle a | D^*b \rangle$$

for all $a, b, c \in A$ and $m \in \mathbb{Z}$.

The radical $\text{Rad}\langle \cdot | \cdot \rangle = \{a \in A \mid \langle a | b \rangle = 0 \ \forall b \in A\}$ of an invariant form is an ideal of A . Also, we have

$$\langle \delta a | b \rangle = \frac{1}{2} \langle [D^*, D]a | b \rangle = \frac{1}{2} \langle a | [D^*, D]b \rangle = \langle a | \delta b \rangle,$$

and this implies that $\langle A_i | A_j \rangle = 0$ for $i \neq j$.

When A is a highest weight module over affine or Virasoro Lie algebra $\mathfrak{g}$ (see e.g. [8, 11]), then one can choose D^* so that the involution $*$ of §2 is the extension of the Chevalley involution of $\mathfrak{g}$, and the contravariant form on A (see e.g. [10]) will be invariant in the sense of Definition 3. See also the example in §5.1.

Remark. Assume that A has an invariant bilinear form $\langle \cdot | \cdot \rangle$. Viewing A as a module over itself, consider the contragredient module A' , see the Remark following Proposition 2 in §2. Then there is an A -algebra homomorphism $\nu : A \rightarrow A'$ given by $a \mapsto \langle a | \cdot \rangle$. In particular, if $\langle \cdot | \cdot \rangle$ is non-degenerate, and all the homogeneous components of A are finite dimensional, then $\nu : A \rightarrow A'$ is an isomorphism. This is a key observation in [12].

Given an invariant form $\langle \cdot | \cdot \rangle$ on A , one can consider a linear map $f : A_0 \rightarrow V$ defined by $f(a) = \langle \mathbf{1} | a \rangle$. Clearly, $f(D^*A_1) = 0$, so that $f \in \text{Hom}(A_0/D^*A_1, V)$, and f defines the form uniquely. We show that this in fact gives an isomorphism of the space of all invariant bilinear forms with $\text{Hom}(A_0/D^*A_1, V)$.

Theorem 1. *Every invariant bilinear form on A is symmetric and there is a one-to-one correspondence between linear maps $f : A_0/D^*A_1 \rightarrow V$ and V -valued invariant bilinear forms $\langle \cdot | \cdot \rangle$ on A , given by $f(a) = \langle \mathbf{1} | a \rangle$ for $a \in A_0$.*

For the proof we need the following lemma.

Lemma 1. *For any $a, b \in A$, if either $m \geq 0$ or $m < -\text{ord } b - 1$, then $a(m)^*b \in D^*A$.*

Proof. Set $\deg a = d$ and $\text{ord } b = k$. Denote $((D^*)^j a)(l) = g_{2(d-j)-l-1}^{d-j-l-1} \in U(A)$, so that

$$g_m^n = ((D^*)^{d+n-m} a)(m-2n-1).$$

For an operator $u \in U(A)$ write $u \sim 0$ if $ub \in D^*A$. Then by (4) we have

$$m g_m^n + g_{m-1}^n + g_{m+1}^{n+1} D^* \sim 0.$$

Consider the generating function $g(x, y) = \sum_{m, n \in \mathbb{Z}} g_m^n x^m y^n$. Let $R : U(A) \rightarrow U(A)$ be the operator given by $Ru = uD^*$. Note that $R^i \sim 0$ for $i > k$, since $(D^*)^{k+1}b = 0$. Then the above relation reads as

$$x \partial_x g + xg + \frac{R}{xy} g \sim 0,$$

therefore

$$g(x, y) \sim \exp\left(-x + \frac{R}{xy}\right) g_0(y)$$

for some series $g_0(y) \in U(A)[[y^{\pm 1}]]$. On the other hand,

$$a(m)^* = (-1)^d \sum_{i \geq 0} (D^{*(i)} a)(2d - m - 2 - i) = (-1)^d \sum_{i \geq 0} \frac{1}{i!} g_{m+1-i}^{m+1-d}.$$

Set $h_m^n = \sum_{i \geq 0} \frac{1}{i!} g_{m-i}^n$, so that $a(m)^* = (-1)^d h_{m+1}^{m+1-d}$. The generating function for h_m^n 's is

$$h(x, y) = \sum_{m, n \in \mathbb{Z}} h_m^n x^m y^n = e^x g(x, y) \sim \exp\left(\frac{R}{xy}\right) g_0(y).$$

Since $(R)^i \sim 0$ for $i > k$, the only powers of x with coefficients $\not\sim 0$ in $h(x, y)$ are $x^0, x^{-1}, \dots, x^{-k}$, therefore $h_m^n \sim 0$ if either $m > 0$ or $m < -k$, and the lemma follows. $\square$

Remark. As a corollary, we get a proof of the fact that $A_d = D^*A_{d+1}$ different from that of Proposition 1. Indeed, take an element $a \in A_d$, $d < 0$. Write $a = a(-1)\mathbf{1}$. Then $a(-1)$ is the dual of $\pm \sum_{i \geq 0} (D^{*(i)} a)(2d - i - 1)$. But $\text{ord } \mathbf{1} = 0$ and $2d - i - 1 \neq -1$ for all $i \geq 0$, so Lemma 1 implies that $a(-1)\mathbf{1} \in D^*A_{d+1}$.

Proof of Theorem 1. We need to show that for every linear map $f : A_0/D^*A_1 \rightarrow V$ there is a symmetric invariant bilinear form $\langle \cdot | \cdot \rangle$ on A such that $f(a) = \langle \mathbf{1} | a \rangle$ for all $a \in A_0$.

For homogeneous $a, b \in A$, define $\langle a | b \rangle = f(a(-1)^*b)$ if $\deg a = \deg b$ and $\langle a | b \rangle = 0$ otherwise. Let us show first that $\langle a(n)b | c \rangle = \langle b | a(n)^*c \rangle$ for all $a, b, c \in A$ and $n \in \mathbb{Z}$. Indeed, it is enough to check that $(a(n)b)(-1)^*c \equiv b(-1)^*a(n)^*c \pmod{D^*A}$. But using the associativity (2), we have $(a(n)b)(-1) - a(n)b(-1) \in U(A)L_+$, and by Lemma 1, $u^*A \subseteq D^*A$ for any $u \in U(A)L_+$. Similarly, $\langle Da | b \rangle = f(a(-1)^*D^*b) = \langle a | D^*b \rangle$. This proves that the form is invariant.

Now we show that the form $\langle \cdot | \cdot \rangle$ is symmetric. We need to have $a(-1)^*b \equiv b(-1)^*a \pmod{D^*A}$ for every $a, b \in A$ of the same degree. Set $u = a(-1)^*b(-1) \in U(A)$ so that $\deg u = 0$. We have to show that

$$(9) \quad (u - u^*)\mathbf{1} \in D^*A_1.$$

By (7), u is a linear combination of operators of the form $c(m)b(-1)$ for some $c \in A$ and $m \in \mathbb{Z}$. But $c(m)b(-1) \equiv (c(m)b)(-1) \pmod{U(A)L_+}$. Hence it is enough to prove (9) for the cases when $u \in U(A)L_+$ and when $u = a(-1)$ for some $a \in A_0$.

If $u \in U(A)L_+$ then $u\mathbb{1} = 0$ and $u^*\mathbb{1} \in D^*A$ by Lemma 1. If $u = a(-1)$ for $a \in A_0$, then

$$a(-1) - a(-1)^* = (a(-1)^* - a(-1))^* = \sum_{i \geq 1} (D^{*(i)}a)(-i-1)^*$$

and again by Lemma 1 we conclude that $(a(-1) - a(-1)^*)\mathbb{1} \in D^*A_1$, since $\text{ord } \mathbb{1} = 0$. $\square$

Remark. It follows from Theorem 1 that if $u \in U(A)$, $\deg u = d$, is such that $u\mathbb{1} = 0$, then $u^*A_d \in D^*A_1$. Indeed, $0 = \langle u\mathbb{1} | A_d \rangle = \langle \mathbb{1} | u^*A_d \rangle$ for every invariant bilinear form on A . In fact, using techniques from [15, 16] one can show that $\{u \in U(A) \mid u\mathbb{1} = 0\} = U(A)L_+$.

4. RADICAL OF A VERTEX ALGEBRA

4.1. The ideal generated by D^*A_1 . Let A be a vertex algebra with an sl_2 -structure as before. Let $I \subset A$ be the ideal generated by D^*A_1 . Denote as usual $I_d = I \cap A_d$.

Proposition 3. (a) $I_d = A_d$ for $d < 0$.

(b) $I_0 = A(-1)D^*A_1$.

(c) A_0/I_0 is an associative commutative algebra with respect to the operation $a \otimes b \mapsto ab = a(-1)b$, and $\mathbb{1}$ is its unit.

Denote $E = A(-1)D^*A_1$. For the proof of Proposition 3 we need the following lemma.

Lemma 2. Let $a \in A_i$ and $b \in A_j$ for $i, j \leq 0$, and if $j = 0$, then $b \in D^*A_1$. Then $a(i+j-1)b \in E$.

Proof. By Proposition 1 we have $a = (D^*)^i a_0$ and $b = (D^*)^j b_0$ for some $a_0 \in A_0$ and $b_0 \in D^*A_1$. Applying (4) $-i$ times we get that

$$a(i+j-1)b \equiv (-1)^i a_0(i+j-1)(D^*)^{-i-j} b_0 \pmod{E}.$$

If $d = i+j = 0$, then $a(i+j-1)b = a(-1)b \in E$ by the definition. We will show by induction on d that $a(d-1)(D^*)^{-d}b \in E$ for any $a \in A_0$, $b \in D^*A_1$ and $d < 0$.

So assume that $a(k-1)(D^*)^{-k}b \in E$ for all $0 \geq k > d$. Then, using (1), we get

$$E \supset D^*A_1 \ni Da(d)(D^*)^{-d}b = a(d)D(D^*)^{-d}b - da(d-1)(D^*)^{-d}b.$$

By Proposition 1, $D(D^*)^{-d}b = (D^*)^{1-d}b_1$ for some $b_1 \in D^*A_1$, so by induction, $a(d)D(D^*)^{-d}b \in E$, and hence also $a(d-1)(D^*)^{-d}b \in E$. $\square$

Remark. In fact, Lemma 2, combined with Lemma 1, shows that $E = U(A)_0 D^*A_1$.

Proof of Proposition 3. (a) follows immediately from Proposition 1.

(b) Let $f : A_0 \rightarrow A_0/E$ be the canonical projection, and let $R = \text{Rad}\langle \cdot | \cdot \rangle$ be the radical of the corresponding form on A . We claim that $R_0 = R \cap A_0 = E$. Indeed, if $a \in A_0 \setminus E$, then $f(a) = \langle a | \mathbb{1} \rangle \neq 0$ and therefore $a \notin R$. In the other direction, take $a \in D^*A_1$. Then for an arbitrary $b \in A_0$ we have

$$\langle a | b \rangle = f(b(-1)^*a) = \sum_{i \geq 0} f((D^{*(i)}b)(-1-i)a) = 0,$$

since $(D^{*(i)}b)(-1-i)a \in E$ by Lemma 2. Therefore, $D^*A_1 \subset R$, and since R is an ideal, we must have $E \subset R$. It follows that $I \subset R_0 = E$, on the other hand obviously $E \in I$, and that proves (b).

(c) Using (2) and Lemma 2 we get

$$(a(-1)b)(-1)c = \sum_{s \geq 0} a(-1-s)b(-1+s)c + \sum_{s \geq 0} b(-2-s)a(s)c \equiv a(-1)b(-1)c \pmod{I_0}$$

for any $a, b, c \in A_0$. By (3) we have $a(-1)b \equiv b(-1)a \pmod{DA_{-1}}$, and we know that $DA_{-1} \subset D^*A_{i+1} \subset I_0 = E$. So the statement (c) follows. $\square$

4.2. The radical.

Definition 4. The radical $\text{Rad } A$ of a vertex algebra A is the radical $\text{Rad}\langle \cdot | \cdot \rangle$ of the invariant bilinear form $\langle \cdot | \cdot \rangle$ corresponding to the projection $f : A_0 \rightarrow A_0/A_0(-1)D^*A_1$.

Remark. This definition has nothing to do with the radical defined in [4].

Recall that by Proposition 3(b), the space $I_0 = A_0(-1)D^*A_1$ is the degree 0 component of the ideal $I \subset A$ generated by D^*A_1 . From the proof of Proposition 3(b) it follows that $(\text{Rad } A)_0 = I_0$. Moreover, the following is true:

Proposition 4. (a) *The radical $R = \text{Rad } A$ is the maximal among all the ideals $J \subset A$ such that $J_0 = A_0(-1)D^*A_1$.*
 (b) $\text{Rad}(A/R) = 0$.

Proof. Let $J \subset A$ be an ideal such that $J_0 = A_0(-1)D^*A_1$. Then for any $a \in J_d$ and $b \in A_d$ we get $\langle a | b \rangle = f(b(-1)^*a) = 0$, therefore $a \in R$. This proves (a). The statement (b) follows from the fact that the form $\langle \cdot | \cdot \rangle$ becomes non-degenerate on A/R . $\square$

4.3. Radical-free algebras. Assume that $R = \text{Rad } A \subsetneq A$. We would like to investigate the quotient vertex algebra A/R . By Proposition 4(b) we have $\text{Rad } A/R = 0$, so we can assume that $\text{Rad } A = 0$. Then $D^*A_1 = I = 0$, the projection $f : A_0 \rightarrow A_0/I_0$ is the identity map, and the corresponding canonical invariant A_0 -valued bilinear form $\langle \cdot | \cdot \rangle$ is non-degenerate.

The following theorem summarizes the properties of radical-free algebras.

Theorem 2. *Let A be a vertex algebra such that $\text{Rad } A = 0$.*

- (a) $A_d = 0$ for $d < 0$.
- (b) A_0 is an associative commutative unital algebra with respect to the product $a \otimes b \mapsto ab = a(-1)b$, with unit $\mathbb{1}$.
- (c) A is a vertex algebra over the associative commutative algebra A_0 .
- (d) Let $I \subset A$ be an ideal. Then $I \cap A_0 = 0$ if and only if $I = 0$.
- (e) For any ideal $J_0 \subset A_0$ the space J_0A is an ideal of A such that $J_0A \cap A_0 = J_0$. It follows that A is simple if and only if A_0 is a field.

Note that all the definitions in §1 make sense when $\mathbb{k}$ is a commutative ring containing $\mathbb{Q}$. So the statement (c) means that elements from $A_0 \subset A$ behave like constants, i.e. A_0 acts on each component A_d by $ab = a(-1)b$ for $a \in A_0$, $b \in A_d$, all structural maps $a(m), D, D^* : A \rightarrow A$ are A_0 -linear, and therefore $DA_0 = 0$ and $A_0(n)A_d = 0$ for $n \neq -1$. Also, the canonical non-degenerate A_0 -valued form $\langle \cdot | \cdot \rangle$ on A is A_0 -bilinear.

Recall that a vertex algebra with sl_2 -structure is called simple if it does not have any D^* -invariant ideals.

Proof. The statements (a) and (b) follow from Proposition 3(a) and (c) respectively, while (d) follows from Proposition 4(a).

Since $D^*A_1 = 0$, we have $\langle Da | b \rangle = 0$ for any $a \in A_0$, $b \in A_1$, therefore $DA_0 \subset \text{Rad } A = 0$. But then $a(n) = 0$ for $n \neq -1$ as an element in the Lie algebra $\text{Coeff } A$, see §1.3. Now the statement (c) follows easily from the formulas (1), (2) and (4) in §1.1.

Finally, I_0A is an ideal of A by (c), and therefore, if A is simple, then A_0 must be a field. This proves one direction of (e). The other direction follows from (d). $\square$

Remark. Let A be a vertex algebra with $\text{Rad } A = 0$ as above. Since $DA_0 = 0$, the formulas (2) and (3) imply that A_1 is a Lie algebra with the bracket $[a, b] = a(0)b$. The bilinear form $\langle \cdot | \cdot \rangle$ on A is an invariant symmetric A_0 -bilinear form on that Lie algebra, and we have $\langle a | b \rangle = a(1)b$ for any $a, b \in A_1$. The Lie algebra A_1 acts on A by derivations of degree 0, so that $ab = a(0)b$ for $a \in A_1$ and $b \in A$. If A is simple, then $A_1(0)A = 0$, since $A_1(0)A$ is always an ideal of A and $\mathbb{1} \notin A_1(0)A$.

If $\mathbb{k} = \mathbb{C}$ and A is a simple radical-free vertex algebra, generated by countably many elements, then we must have $A_0 = \mathbb{C}$. We see that radical-free vertex algebras are very close to be of a CFT type, after we set $\mathbb{k} = A_0$. The only thing that might be missing is the condition $\dim A_d < \infty$.

Question. Are there any radical-free vertex algebras A with sl_2 -structure, such that the homogeneous components A_d have infinite rank over A_0 ?

5. EXAMPLES

5.1. Heisenberg algebra. Let A be the Heisenberg vertex algebra. By definition, it is generated by a single element $a \in A_1$ such that the locality $N(a, a) = 2$ and $a(0)a = 0$, $a(1)a = \mathbb{1}$. One can show (see e.g. [16]) that these conditions define A uniquely. We have $A_0 = \mathbb{k}\mathbb{1}$ and $A_1 = \mathbb{k}a$. It is well known, that A is a simple vertex algebra in the sense that it does not have any ideals at all, regardless of any sl_2 -structure.

For every $k \in \mathbb{k}$ the element $\omega_k = \frac{1}{2}a(-1)a + kDa \in A_2$ is a Virasoro element of A . If we set $D^* = \omega_0(2)$, then $D^*a = 0$, hence $A_0/D^*A_1 = \mathbb{k}\mathbb{1}$, and therefore A has only one invariant bilinear form $\langle \cdot | \cdot \rangle$, up to a scaling factor, and in this case $\text{Rad } A = \text{Rad} \langle \cdot | \cdot \rangle = 0$. If instead we set $D^* = \omega_k(2)$ for $k \neq 0$, then $D^*a = -2k\mathbb{1}$. In this case $A_0/D^*A_1 = 0$, $\text{Rad } A = A$, and A does not have any invariant bilinear forms.

5.2. Free vertex algebras. Let $\mathcal{G}$ be a set and $N : \mathcal{G} \times \mathcal{G} \rightarrow \mathbb{Z}$ be a symmetric function such that $N(a, a) \in 2\mathbb{Z}$. In [16] we have constructed a vertex algebra $F = F_N(\mathcal{G})$ which we called *the free vertex algebra generated by $\mathcal{G}$ with respect to locality bound N* . It is generated by $\mathcal{G}$ so that the locality of any $a, b \in \mathcal{G}$ is exactly $N(a, b)$, and for any other vertex algebra A generated by $\mathcal{G}$ with the same bound on locality, there is a homomorphism $F \rightarrow A$ preserving $\mathcal{G}$. The algebra F is graded by the semilattice $\mathbb{Z}_+[\mathcal{G}]$, we refer to this grading as the grading by weights. Also, F is graded by degrees if we set $\deg a = -\frac{1}{2}N(a, a)$ for $a \in \mathcal{G}$. If we denote by $F_{\lambda, d}$ the space of all elements of F of weight $\lambda \in \mathbb{Z}_+[\mathcal{G}]$ and degree $d \in \mathbb{Z}$, then

$$F_{\lambda, k}(n)F_{\mu, l} \subseteq F_{\lambda+\mu, k+l-n-1}, \quad DF_{\lambda, k} \subseteq F_{\lambda, k-1} \quad \text{and} \quad \mathbb{1} \in F_{0, 0}.$$

It is not difficult to see that we can define an operator $D^* : F \rightarrow F$, satisfying (4), by setting $D^*a = 0$ for all $a \in \mathcal{G}$. Then we have $D^*F_{\lambda, d} \subseteq F_{\lambda, d+1}$.

It is shown in [16] that for any weight $\lambda = a_1 + \dots + a_l \in \mathbb{Z}_+[\mathcal{G}]$ there is the minimal degree

$$d_{\min}(\lambda) = -\frac{1}{2} \sum_{i, j=1}^l N(a_i, a_j) \in \mathbb{Z}$$

such that $\dim_{\mathbb{k}} F_{\lambda, d_{\min}(\lambda)} = 1$ and $\dim_{\mathbb{k}} F_{\lambda, d} = 0$ for $d < d_{\min}(\lambda)$. For $d > d_{\min}(\lambda)$ the dimension of $F_{\lambda, d}$ is equal to the number of partitions of $d - d_{\min}(\lambda)$ into a sum of l non-negative integers colored by $a_1, \dots, a_l$.

It is also proved in [16] that F is embedded into the vertex algebra V_{Λ} corresponding to the lattice $\Lambda = \mathbb{Z}[\mathcal{G}]$. The scalar product on Λ is defined by $\langle a | b \rangle = -N(a, b)$ for $a, b \in \mathcal{G}$.

Since D^* is homogeneous with respect to both weights and degrees, and the only element of weight 0 in F is $\mathbb{1}$, up to a scalar, it follows that $\mathbb{1} \notin D^*F_1$. So one can always define a functional $f : F_0/D^*F_1 \rightarrow \mathbb{k}$ so that $f(\mathbb{1}) = 1$ and $f(F_{\lambda, 0}) = 0$ for every weight $\lambda \neq 0$. The corresponding invariant bilinear form on F is a restriction of the canonical bilinear form on V_{Λ} . If $N(\mathcal{G}, \mathcal{G}) < 0$, then this will be the only form on F , up to a scalar. If, on the contrary, the localities $N(a_i, a_j)$ are large, then $\dim F_{\lambda, 0}$ grows as λ gets longer, and it follows from the results of [16] that $\dim F_{\lambda, 0}/(\text{Rad } F)_{\lambda, 0} > 0$ when $\lambda \in \mathbb{Z}_+[\mathcal{G}]$ is big enough. Hence, in this case there are infinitely many different invariant bilinear forms on F , and infinitely many simple radical-free vertex algebras, which are homomorphic images of F .

Remark. It appears that the algebra $\bar{F} = F/\text{Rad } F$ is a very interesting object from combinatorial point of view. It is possible to show that the commutative associative algebra $\bar{F}_0$ is isomorphic to a polynomial algebra in infinitely many variables. Also, the Question at the end of §4 is equivalent to asking whether the components $\bar{F}_d$ have finite rank as modules over $\bar{F}_0$.

Acknowledgments. I thank Haisheng Li for inspiring discussions and for pointing out an inaccuracy in an earlier version of this paper.

REFERENCES

- [1] R. E. Borcherds. Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Nat. Acad. Sci. U.S.A.*, 83(10):3068–3071, 1986.
- [2] C. Dong. Vertex algebras associated with even lattices. *J. Algebra*, 161(1):245–265, 1993.
- [3] C. Dong, H. Li, and G. Mason. Vertex Lie algebras, vertex Poisson algebras and vertex algebras. In *Recent developments in infinite-dimensional Lie algebras and conformal field theory (Charlottesville, VA, 2000)*, volume 297 of *Contemp. Math.*, pages 69–96. Amer. Math. Soc., Providence, RI, 2002.
- [4] C. Dong, H. Li, G. Mason, and P. S. Montague. The radical of a vertex operator algebra. In *The Monster and Lie algebras (Columbus, OH, 1996)*, volume 7 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 17–25. de Gruyter, Berlin, 1998.
- [5] E. Frenkel and D. Ben-Zvi. *Vertex algebras and algebraic curves*, volume 88 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2001.
- [6] I. B. Frenkel, Y.-Z. Huang, and J. Lepowsky. On axiomatic approaches to vertex operator algebras and modules. *Mem. Amer. Math. Soc.*, 104(494), 1993.
- [7] I. B. Frenkel, J. Lepowsky, and A. Meurman. *Vertex Operator Algebras and the Monster*. Academic Press, Boston, MA, 1988.
- [8] I. B. Frenkel and Y. Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, 66(1):123–168, 1992.
- [9] R. L. Griess Jr. GNAVOA, I, 2001. To appear in Proceedings of the Infinite Dimensional Lie Theory Meeting, Fields Institute, 23-27 October, 2000.
- [10] V. G. Kac. *Infinite-Dimensional Lie Algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [11] V. G. Kac. *Vertex Algebras for Beginners*, volume 10 of *University Lecture Series*. AMS, Providence, RI, second edition, 1998.
- [12] H. Li. Symmetric invariant bilinear forms on vertex operator algebras. *J. Pure Appl. Algebra*, 96(3):279–297, 1994.
- [13] H. Li. An analogue of the Hom functor and a generalized nuclear democracy theorem. *Duke Math. J.*, 93(1):73–114, 1998.
- [14] M. Primc. Vertex algebras generated by Lie algebras. *J. Pure Appl. Algebra*, 135(3):253–293, 1999.
- [15] M. Roitman. On free conformal and vertex algebras. *J. Algebra*, 217(2):496–527, 1999.
- [16] M. Roitman. Combinatorics of free vertex algebras. *J. Algebra*, 255(2):297–323, 2002.
- [17] M. Roitman. On Griess algebras, 2003. In preparation.

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, ANN ARBOR, MI 48109-1109

E-mail address: roitman@umich.edu